

# Some slides for 16th Lecture, Algebra 2

Diego Ruano

Department of Mathematical Sciences  
Aalborg University  
Denmark

14-04-2014

## Lemma 4.8.1

Let  $F$  be a finite field. Then  $|F| = p^n$ , where  $p$  is a prime number,  $n \geq 1$  and there exists an irreducible polynomial  $f \in \mathbb{F}_p[X]$  of degree  $n$  such that

$$F \cong \mathbb{F}_p[X] / \langle f \rangle$$

### Theorem 4.8.2

There exists a unique finite field with  $p^n$  elements, where  $p$  is a prime number and  $n \geq 1$ . More precisely, we have

- 1 There exists an irreducible polynomial in  $\mathbb{F}_p[X]$  of degree  $n$ .
- 2 Suppose that  $F$  and  $F'$  are finite fields with  $p^n$  elements. Then there exists a ring isomorphism  $F \cong F'$ .

### Lemma 4.8.3

Let  $\tau$ ,  $d$  and  $n$  be natural numbers, where  $\tau > 1$ . Then  $\tau^d - 1$  divides  $\tau^n - 1$  if and only if  $d$  divides  $n$ .

## Theorem 4.8.5

There exists an irreducible polynomial in  $\mathbb{F}_p[X]$  of degree  $n \geq 1$ . More precisely, if  $f$  is an irreducible polynomial dividing  $\Phi_{p^n-1}$  in  $\mathbb{F}_p[X]$  then  $\deg(f) = n$ .

# Uniqueness of finite fields

